

## **Publications from our previous Massachusetts Hispanic Elders Study (MAHES)**

Merete C, Falcón LM, Tucker KL. Vitamin B6 is associated with Depression in Massachusetts elders. *J Am Coll Nutr.* 2008;27: 421-7.

Gao X, Nelson ME, Tucker KL. Television viewing is associated with prevalence of metabolic syndrome in Hispanic elders. *Diabetes Care.* 2007;30:694–700.

Seaverson EL, Buell J, Fleming DJ, Bermudez O, Wood R, Potischman N, Chasan-Taber L, Tucker KL. Poor iron status is more prevalent in Hispanic than in non-Hispanic white older adults in Massachusetts. *J Nutr.* 2007;137:414-20.

Gao X, Martin A, Lin H, Bermudez OI, Tucker KL. Alpha-tocopherol intake and plasma concentration of Hispanic and non-Hispanic white elders is associated with dietary intake pattern. *J Nutr.* 2006;136:2574-9.

Bermudez OI, Ribaya-Mercado JD, Talegawkar SA, Tucker KL. Hispanic and non-Hispanic white elders from Massachusetts have different patterns of carotenoid intake and plasma concentrations. *J Nutr.* 2005;135:1496-1502.

Gao X, Bermudez OI, Tucker KL. Plasma C-reactive protein and homocysteine concentrations are related to frequent fruit and vegetable intake in Hispanic elders. *J Nutr.* 2004;134:913-8.

Lin H, Bermudez OI, Tucker KL. Dietary patterns of Hispanic elders are associated with acculturation and obesity. *J Nutr.* 2003;133:3651-7.

Bermudez OI, Velez-Carrasco W, Schaefer EJ, Tucker KL. Dietary and plasma lipids, lipoprotein and apolipoprotein profiles among elderly Hispanics and non-Hispanics and their association with diabetes. *Am J Clin Nutr.* 2002;76:1214-21.

Lin H, Bermudez OI, Falcon LM, Tucker KL. Hypertension among Hispanic elders in Massachusetts. *Ethnic & Disease.* 2002;12:499-507.

Kwan L, Bermudez OI, Tucker KL. Low vitamin B12 intake and status are more prevalent in Hispanic older adults of Caribbean origin than in neighborhood-matched non-Hispanic Whites. *J Nutr.* 2002;132:2059-64.

Chen H, Bermudez OI, Tucker KL. Waist circumference and weight change are associated with disability among elderly Hispanics. *J Gerontol: Med Sci.* 2002;57:M19-M25.

Bermudez OI, Tucker KL. Total and Central Obesity among Elderly Hispanics and the Association with Type 2 Diabetes. *Obesity Res.* 2001;9(8):443-51.

Tucker KL, Bermudez O, Castaneda C. Type 2 diabetes is prevalent and poorly controlled among Hispanic elders. *Am J Pub Health*. 2000;90:1288-93.

Bermudez OI, Tucker KL. Uso de la altura de rodilla para corregir la talla de ancianos de origen Hispano. *Archivos Latinoamericanos de Nutricion*. 2000;50(1):42-7.

Bermudez OI, Falcon LM, Tucker KL. Intake and food sources of macronutrients among Hispanic elders: Association with ethnicity, acculturation and length of residence in the United States. *J Am Dietet Assoc*. 2000;100:665-73.

Castaneda C, Bermudez O, Tucker KL. Protein nutritional status and function are associated with Type 2 diabetes in Hispanic elders. *Am J Clin Nutr*. 2000;72:89-95.

Falcon LM, Tucker KL. Prevalence and correlates of depressive symptomatology among Hispanic elders in Massachusetts. *J Gerontol: Soc Sci*. 2000;55B(2):S108-S116.

Tucker KL, Falcon LM, Bianchi LI, Cacho E, Bermudez OI. Self reported prevalence and health correlates of functional limitation among Massachusetts elderly Puerto Ricans, Dominicans, and a non-Hispanic white neighborhood comparison group. *J Gerontol: Med Sci*. 2000;55A(2):M90-M97.

Bermudez O, Becker E, Tucker KL. Development of sex-specific equations for estimating stature of frail elderly Hispanics living in the Northeastern United States. *Am J Clin Nutr*. 1999;69:992-8.

Tucker KL, Bianchi L, Maras J, and Bermudez O. Adaptation of a food frequency questionnaire to assess diets of Puerto Rican and non-Hispanic adults. *Am J Epidemiol*. 1998;148:507-18.